

Stability by KAM confinement of certain wild, nongeneric relative equilibria of underwater vehicles with coincident centers of mass and bouyancy

George W. Patrick

April 2002

Abstract

Purely rotational relative equilibria of an ellipsoidal underwater vehicle occur at nongeneric momentum where the symplectic reduced spaces change dimension. The stability these relative equilibria under momentum changing perturbations is not accessible by Lyapunov functions obtained from energy and momentum. A blow-up construction transforms the stability problem to the analysis symmetry-breaking perturbations of Hamiltonian relative equilibria. As such, the stability follows by KAM theory rather than energy-momentum confinement.

INTRODUCTION

The phase space $TSE(3)$ with Lagrangian

$$L(A, a, \Omega, v) \equiv \frac{1}{2} \Omega^t \mathbf{I} \Omega + \frac{1}{2} v^t \mathbf{M} v \quad (1)$$

approximately models the motion of a neutrally buoyant vehicle submerged in an inviscid irrotational fluid (see Leonard [1997] and the references therein), in the case of coincident centers of mass and buoyancy. Here tangent vectors of $SE(3)$ are represented by left translation and elements of $SE(3)$ parameterize the configurations of the vehicle by embedding a reference vehicle into the fluid. $\mathbf{I}$ and $\mathbf{M}$ are constant, positive definite, 3×3 matrices that can be calculated from the shape and mass distribution of the vehicle. This system admits the $SE(3)$ symmetry of the left action of $SE(3)$ on itself.

For an ellipsoidal vehicle with principle axes of inertia along the axes of symmetry of the ellipsoid, $\mathbf{I} = \text{diag}(I_1, I_2, I_3)$ and $\mathbf{M} = \text{diag}(M_1, M_2, M_3)$ (i.e. $\mathbf{I}$ and $\mathbf{M}$ are diagonal). If $M_1 = M_2$ and $I_1 = I_2$ (or similarly if $M_1 = M_3$ and $I_1 = I_3$ etc.) then there is a further material symmetry of the system: $SO(2) = \{ \exp(\mathbf{k} \wedge \theta) \}$ acts as a subgroup of $SE(3)$ by inverse multiplication on

the right. In the case that $\mathbf{I}$ and $\mathbf{M}$ are both constants of the identity then the material symmetry is $SO(3)$.

Lie-Poisson reduction yields the Poisson phase space $\mathfrak{se}(3)^* = \{(\pi, p)\}$ where $\pi = \mathbf{I}\Omega$ and $p = \mathbf{M}v$. The equations of motion are

$$\frac{d\pi}{dt} = \pi \times \Omega + p \times v, \quad \frac{dp}{dt} = p \times \Omega, \quad (2)$$

and by direct substitution, for each $\alpha_e \in \mathbb{R}$,

$$p_e^{\alpha_e} : \quad \pi = \alpha_e \mathbf{k}, \quad p = 0,$$

is an equilibrium of the Poisson reduced systems and hence a relative equilibrium of the original system. The generator is

$$\Omega_e^{\alpha_e} \equiv \frac{\alpha_e}{I_3} \mathbf{k}, \quad v_e^{\alpha_e} \equiv 0,$$

so the relative equilibrium corresponds to a stationary vehicle rotating about an axis of symmetry which is aligned with the vertical. This article is concerned with the stability of these relative equilibria in the case where I_3 is not an intermediate axis of symmetry, i.e. assuming $I_1 < I_2 < I_3$ or $I_3 < I_2 < I_1$.

The symplectic leaves of $se(3)^*$ are as follows. On the complement of $p = 0$ lie the generic symplectic leaves, all diffeomorphic to TS^2 , and which are the level sets of the two Casimirs $|p|$ and $\pi \cdot p$. Nongeneric leaves occur within the set $p = 0$ and are the level sets of the subcasimir $|\pi|$. Thus the relative equilibria $p_e^{\alpha_e}$ correspond to Lyapunov stable equilibria on the (nongeneric) symplectic leaves of $se(3)^*$ since the energy has a definite critical point when restricted to those leaves. Were the symmetry group to be compact this *leafwise stability* would imply stability of the equilibrium modulo the isotropy group of the momentum. $SE(3)$, of course, is not compact. The question is whether or not $p_e^{\alpha_e}$ are stable under perturbations from nongeneric leaves into nearby generic leaves.

Leonard and Marsden [1997] have identified this question as particularly delicate, and the theory of Patrick, Roberts, and Wulff [2002], the sharpest possible for the problem of establishing the stability of relative equilibria by energy-momentum confinement in the case of noncompact symmetry, corroborates that opinion. Patrick *et al.* separate generators of relative equilibria into two complementary classes, *tame* and *wild*. The generator of $p_e^{\alpha_e}$ is tame if and only if $\alpha_e = 0$, corresponding to a stationary, nonrotating vehicle, in which case $SE(3)$ -stability follows directly, since the energy has zero derivative at $p_e^{\alpha_e}$ and has positive definite Hessian there. However, if $\alpha_e \neq 0$, the generator is wild and the theory does not imply stability.

So it is an open question whether the relative equilibria $p_e^{\alpha_e}$, $\alpha_e \neq 0$, are $SE(3)$ -stable or not, and the problem appears inaccessible by energy-momentum confinement. This is due to the presence of a noncompact symmetry group and wild generators.

§1 THE BLOW-UP CONSTRUCTION

In its essence the stability issue is one of perturbation from a nongeneric symplectic leaf to nearby, higher dimensional, generic leaves. In order to bridge to Hamiltonian perturbation theory, which is usually cast in a setting of a fixed canonical phase space, it is natural to begin by normalizing the generic leaves. The leaf corresponding to

$$|p| = a, \quad \pi \cdot p = b,$$

for $a > 0$ is diffeomorphic to

$$TS^2 = \{(w, \dot{w}) \in \mathbb{R}^3 \times \mathbb{R}^3 : |w| = 1, w \cdot \dot{w} = 0\}$$

by the map

$$w = \frac{p}{|p|}, \quad \dot{w} = \pi - \frac{\pi \cdot p}{|p|^2} p,$$

the inverse map being given by, for fixed $a > 0$ and $b \in \mathbb{R}$,

$$p = aw, \quad \pi = \dot{w} + \frac{b}{a} w. \tag{3}$$

Having normalized the symplectic leaves to the constant manifold TS^2 , one seeks to extend this to the nongeneric leaves within $p = 0$, which means extending it to $a = 0$, since $p = aw$. As it stands (3) is poorly defined for $a = 0$, but for fixed ratios of b/a it is well defined even for arbitrarily small a , suggesting that the proper way to approach the nongeneric leaves from generic ones is through constant $\pi \cdot p/|p|$. *Setting $\gamma \equiv b/a$ and using the parameters a and γ instead of a and b codes the generic leaves so they fit smoothly into the nongeneric ones, thus allowing the possibility of an effective perturbation approach.* The map $p = aw$, $\pi = \dot{w} + \gamma w$ for $a = 0$ is many-to-one and so the three dimensional set of nongeneric leaves $p = 0$ is “blown-up” by this map to the five dimensional set of $TS^2 \times \mathbb{R} = \{(w, \dot{w}), \gamma\}$. Thus one is led to define the *blown-up space* of $se(3)^*$ as

$$\hat{P} \equiv TS^2 \times \mathbb{R}_{\geq 0} \times \mathbb{R} \equiv \{(w, \dot{w}, a, \gamma) : |w| = 1, w \cdot \dot{w} = 0, a \geq 0\}$$

with *blow-down map*

$$p = aw, \quad \pi = \dot{w} + \gamma w$$

and corresponding *blow-up map*, defined on the generic ($p \neq 0$) leaves only,

$$w = \frac{p}{|p|}, \quad \dot{w} = \pi - \frac{\pi \cdot p}{|p|^2} p, \quad a = |p|, \quad \gamma = \frac{b}{a}.$$

The blow-up map is a diffeomorphism from the (open) set of generic leaves to the (open) set $a > 0$ in the blown-up space (the *generic sector*), such that each

generic leaf is sent to the constant manifold TS^2 . The evolution of the generic leaves is transformed to a evolution on TS^2 parameterized by the Casimir values a and γ . The blow-down map takes the set $a = 0$ in the blown-up space (the *nongeneric sector*) to the set of nongeneric leaves, and is many-to-one on that sector. Increasing the parameter a from zero corresponds to leaving the nongeneric leaves and moving to the generic ones, while γ parameterizes the possible avenues of departure.

The utility of the blow-up to support perturbation arguments depends on whether or not the dynamics of the generic sector can be continued smoothly to the nongeneric sector. On the generic sector the vector field that generates the dynamics is

$$\begin{aligned}
\frac{dw}{dt} &= \frac{1}{a} \frac{dp}{dt} = \frac{1}{a} p \times \mathbf{I}^{-1} \pi = w \times \mathbf{I}^{-1} (\dot{w} + \gamma w), \\
\frac{d\dot{w}}{dt} &= \frac{d\pi}{dt} - \gamma \frac{dw}{dt} \\
&= \pi \times \mathbf{I}^{-1} \pi + p \times \mathbf{M}^{-1} p - \gamma \frac{dw}{dt} \\
&= (\dot{w} + \gamma w) \times \mathbf{I}^{-1} (\dot{w} + \gamma w) + aw \times \mathbf{M}^{-1} aw - \gamma w \times \mathbf{I}^{-1} (\dot{w} + \gamma w) \\
&= \dot{w} \times \mathbf{I}^{-1} (\dot{w} + \gamma w) + a^2 w \times \mathbf{M}^{-1} w.
\end{aligned} \tag{4}$$

Obviously this is smooth in a for all $a \geq 0$, as required. Dynamics on $a = 0$ that is robust enough to continue through perturbation to small positive a will have implications for the original system. By continuity in a , the blown-up vector field is a lift by the smooth blow-down map of the vector field for the original system, even through the nongeneric sector. Thus the flow on the nongeneric sector corresponds through the blow-down map to the flow of the original system on the union of the nongeneric leaves.

Actually, the blow-up has a very transparent reformulation, since $\hat{P}$ is diffeomorphic to $S^2 \times \mathbb{R}^3 \times \mathbb{R}_{\geq 0} = \{(w, \pi, a)\}$ by the map $\pi = \gamma w + \dot{w}$. Through this diffeomorphism the blow-down map is simply $p = aw$, which is to say that w by itself is enough to desingularize the foliation by symplectic leaves, but not enough to normalize the leaves. The blow up map is a proper map since the map $(a, w) \mapsto aw$ is proper.

Some exploration of the nongeneric sector of the blown-up space may be helpful for visualization purposes. For fixed π_0 the equation $\gamma w + \dot{w} = \pi_0$ has solution $w \in S^2$ and $\gamma = \pi_0 \cdot w$. Consequently the blow-up of the point $p = 0$, $\pi = \pi_0$ is in all cases a two sphere. This sphere intersects fixed γ such that $|\gamma| < |\pi_0|$ in a circle, $\gamma = \pm |\pi_0|$ in a point, and $|\gamma| > |\pi_0|$ not at all. Thus departure from the point $p = 0$, $\pi = \pi_0$ along $\gamma > |\pi_0|$ is impossible. The nongeneric symplectic leaf $|\pi| = r > 0$ blows up to $\gamma^2 + |\dot{w}|^2 = r^2$ which is diffeomorphic to $S^2 \times S^2$, and which for fixed $|\gamma| < r$ is a circle bundle and for $|\gamma| = r$ is a sphere.

The Hamiltonian pulls back through the smooth blow-down map to

$$\begin{aligned}\hat{H} &\equiv \frac{1}{2}\pi^t \mathbf{I}^{-1} \pi + \frac{1}{2}p^t \mathbf{M}^{-1} p = \hat{H}^0 + a^2 \hat{H}^1, \\ \hat{H}^0 &\equiv \frac{1}{2}(\dot{w} + \gamma w)^t \mathbf{I}^{-1} (\dot{w} + \gamma w), \quad \hat{H}^1 \equiv \frac{1}{2}w^t \mathbf{M}^{-1} w,\end{aligned}$$

$\hat{H}$ is written this way in anticipation of perturbation arguments from $a = 0$ to small nonzero a . The symplectic form $\hat{\omega}$ on the nongeneric sector can be calculated from the formula for the coadjoint orbit symplectic forms of $SE(3)$ in Marsden and Ratiu [1994], with the result that

$$\begin{aligned}\hat{\omega}(w, \dot{w})((\delta w_1, \delta \dot{w}_1), (\delta w_2, \delta \dot{w}_2)) \\ = -w \cdot (\delta w_1 \times \delta \dot{w}_2 - \delta w_2 \times \delta \dot{w}_1) - \gamma w \cdot (\delta w_1 \times \delta w_2).\end{aligned}$$

By continuity, the relation $i_{X_{\hat{H}}} \hat{\omega} = d\hat{H}$ persists from $a = 0$ to $a > 0$, so the vector field (4) is Hamiltonian at $a = 0$ with symplectic form $\hat{\omega}$ and Hamiltonian $\hat{H}^0$. Thus the evolution on the nongeneric sector is Hamiltonian, in a way that smoothly continues the Hamiltonian structure of the generic sector.

The dynamics on the invariant submanifold $p = 0$ in the original space P admits the subcasimir $|\pi|$. This conserved quantity (*conserved on $p = 0$ only*) pulls back to a conserved quantity $|w + \gamma \dot{w}|$ for the *nongeneric sector* of the blow-up space $\hat{P}$. Since $w \cdot \dot{w} = 0$ and $|w| = 1$, this gives the conserved quantity $|\dot{w}|^2$ and hence the conserved quantity $f(|\dot{w}|)$, where f is any function. The Hamiltonian vector field of $f(|\dot{w}|^2)$ is

$$\frac{dw}{dt} = -\frac{f'(|\dot{w}|)}{|\dot{w}|} \dot{w} \times w, \quad \frac{d\dot{w}}{dt} = -\gamma \frac{f'(|\dot{w}|)}{|\dot{w}|} w \times \dot{w}.$$

Note that $\tilde{m} \equiv \dot{w} + \gamma w$ is conserved by these equations, so that

$$\frac{dw}{dt} = -\frac{f'(|\dot{w}|)}{|\dot{w}|} \tilde{m} \times w, \quad \frac{d\dot{w}}{dt} = -\gamma \frac{f'(|\dot{w}|)}{|\dot{w}|} \tilde{m} \times \dot{w},$$

the solution of which is rotations about $\tilde{m}$. To normalize the period at 2π and the righthand sense about $\tilde{m}$, choose

$$\frac{f'(|\dot{w}|)}{|\dot{w}|} |\tilde{m}| = \frac{f'(|\dot{w}|)}{|\dot{w}|} \sqrt{\gamma^2 + |\dot{w}|^2} = -1,$$

which gives $f(|\dot{w}|) = -\sqrt{\gamma^2 + |\dot{w}|^2}$. Thus the nongeneric sector has an *additional $SO(2)$ symmetry*, which acts by

$$\theta \cdot (w, \dot{w}) \equiv (\exp(m^\wedge \theta) w, \exp(m^\wedge \theta) \dot{w}), \quad m \equiv \frac{\dot{w} + \gamma w}{\sqrt{\gamma^2 + |\dot{w}|^2}},$$

and has momentum

$$\hat{J} \equiv -\sqrt{\gamma^2 + |\dot{w}|^2}.$$

This extra $SO(2)$ symmetry arises from a subcasimir of the original system. The action and the corresponding momentum are defined on the nongeneric sector where γ and $\dot{w}$ are not both zero. The set where $a = \gamma = 0$ and $\dot{w} = 0$ exactly corresponds through the blow-down/up to the set where $p = 0$ and $\pi = 0$, so that the action and its momentum are defined only on an open subset does not affect the analysis of the relative equilibria $p_e^{\alpha_e}$.

Here are some aspects of the $SO(2)$ action and its relation to the blow-down/up map.

1. The action is free except on the set $\dot{w} = 0$, which is a 2-sphere of fixed points. This two sphere is also the level $-|\gamma|$ of the momentum $\hat{J}$, is a symplectic submanifold of $\hat{P}$, and as such is equal to its own singular reduction.
2. The orbit relation of the action together with the parameter γ exactly absorb the additional phase space from blowing up the nongeneric leaves. Indeed, for fixed γ the blow-down map is a quotient map for the action, and the orbit space is therefore smooth, irrespective of the fact that the action is not free.
3. The blow-down map restricts to a quotient map for the (singular or nonsingular) symplectic reduced space associated to the $\hat{J} = \hat{\mu}$ level set. As such this reduced space is symplectomorphic to the nongeneric leaf $p = 0$, $|\pi| = -\hat{\mu}$.

Only the verification of the third item in the nonsingular ($\hat{\mu} < -|\gamma|$) case is troublesome. For that, it is easily verified that the map $\pi = \gamma w + \dot{w}$ is a quotient map for the $SO(2)$ action on $\hat{J}^{-1}(\hat{\mu})$ which has image the sphere $TS_{-\hat{\mu}}^2 = \{\pi : |\pi| = -\hat{\mu}\}$. To pull down the symplectic form $\hat{\omega}$ by that map, first let $(\pi, \delta\pi_i) \in TS_{-\hat{\mu}}^2$, $i = 1, 2$, and seek $(w, \dot{w}, \delta w_i, \delta \dot{w}_i)$ such that

$$\begin{aligned} |w| &= 1, & w \cdot \dot{w} &= 0, & w \cdot \delta w_i &= 0, & \delta w_i \cdot \dot{w} + w \cdot \delta \dot{w}_i &= 0, \\ -\sqrt{\gamma^2 + |\dot{w}|^2} &= \hat{\mu}, & \delta \dot{w}_i \cdot \dot{w} &= 0, & \pi &= \gamma w + \dot{w}, & \delta \pi_i &= \gamma \delta w_i + \delta \dot{w}_i. \end{aligned}$$

To solve these equations choose w such that $w \cdot \hat{\pi} = \gamma$ and set $\dot{w} = \pi - \gamma w$. Expanding δw_i and $\delta \dot{w}_i$ in the basis $w, \dot{w}, w \times \dot{w}$ gives

$$\delta w_i = -\frac{w \cdot \delta \pi_i}{\hat{\mu}^2 - \gamma^2}(\pi - \gamma w), \quad \delta \dot{w}_i = (w \cdot \delta \pi_i)w + \frac{(w \times \pi) \cdot \delta \pi_i}{\hat{\mu}^2 - \gamma^2}w \times \pi.$$

Substitution into $\hat{\omega}$ then gives

$$\begin{aligned}
& \hat{\omega}(w, \dot{w})((\delta w_1, \delta \dot{w}_1), (\delta w_2, \delta \dot{w}_2)) \\
&= \frac{1}{\hat{\mu}^2 - \gamma^2} \left((\delta \pi_1 \cdot w) ((\pi \times \delta \pi_2) \cdot w) - (\delta \pi_2 \cdot w) ((\pi \times \delta \pi_1) \cdot w) \right) \\
&= \frac{1}{\hat{\mu}^2 - \gamma^2} (w \times (w \times \pi)) \cdot (\delta \pi_1 \times \delta \pi_2) \\
&= \frac{1}{\hat{\mu}^2 - \gamma^2} ((w \cdot \pi)w - \pi) \cdot (\delta \pi_1 \times \delta \pi_2) \\
&= \frac{1}{\hat{\mu}^2 - \gamma^2} \left((w \cdot \pi) \frac{w \cdot \pi}{|\pi|^2} \pi - \pi \right) \cdot (\delta \pi_1 \times \delta \pi_2) \\
&= \frac{1}{\hat{\mu}^2 - \gamma^2} \left(\frac{\gamma^2}{\hat{\mu}^2} - 1 \right) \pi \cdot (\delta \pi_1 \times \delta \pi_2) \\
&= -\frac{1}{|\pi|^2} \pi \cdot (\delta \pi_1 \times \delta \pi_2).
\end{aligned}$$

This last expression is the symplectic form on the nongeneric leaf $|\pi| = -\hat{\mu}$, as required.

In short, the $SO(2)$ symmetry arises and exactly absorbs the additional dimensions resulting from the blow-up construction. The symplectic reductions of the nongeneric sector by this symmetry exactly coincide with the original system restricted to the nongeneric symplectic leaves of the phase space P .

The pull-back of the relative equilibria $p_e^{\alpha_e}$ by the blow-down map is the set of $(w, \dot{w}, a, \gamma)$ such that

$$p = aw = 0, \quad \pi = \alpha_e \mathbf{k} = \dot{w} + \gamma w.$$

Since $|w| = 1$ the first equation is equivalent to $a = 0$ (the relative equilibria are of course in the nongeneric sector), and dotting the second with w shows it is equivalent to $\gamma = \alpha_e \mathbf{k} \cdot w$ and $\dot{w} = \alpha_e \mathbf{k} - \gamma w$. Since $\gamma^2 + |\dot{w}|^2 = \alpha_e^2$ and $\alpha_e \neq 0$, all these solutions are in the open set where the $SO(2)$ action is defined (i.e. where γ and $\dot{w}$ are not both zero). Thus the relative equilibria $p_e^{\alpha_e}$ blow up to

$$\hat{p}_e^{\alpha_e} : \quad |w| = 1, \quad \dot{w} = \alpha_e \mathbf{k} - \alpha_e (\mathbf{k} \cdot w) w, \quad a = 0, \quad \gamma = \alpha_e \mathbf{k} \cdot w. \quad (5)$$

Since w is unconstrained in (5), except for the first equation, $\hat{p}_e^{\alpha_e}$ is diffeomorphic to S^2 . Since $m = \mathbf{k}$ on $\hat{p}_e^{\alpha_e}$, the $SO(2)$ action on $\hat{p}_e^{\alpha_e}$ is by rotation of the pair $(w, \dot{w})$ about $\mathbf{k}$. By substitution of $\hat{p}_e^{\alpha_e}$ into (4), each point of $\hat{p}_e^{\alpha_e}$ is a relative equilibrium for the $SO(2)$ symmetry, except for the two points $w = \pm \mathbf{k}, \dot{w} = 0, a = 0, \gamma = \pm \alpha_e$, which are equilibria that reside at singular points of the action. Each relative equilibrium in $\hat{p}_e^{\alpha_e}$ has the same generator, namely

$$\hat{\xi}_e^{\alpha_e} \equiv -\frac{\alpha_e}{I_3} \quad (6)$$

and by substitution into $\hat{J}$, the same momentum, namely

$$\hat{\mu}_e^{\alpha_e} \equiv -\alpha_e.$$

Fixing γ , which means fixing a parameter, $\alpha_e = \gamma/\cos\phi$, where ϕ is the angle between $\mathbf{k}$ and w . Thus for fixed γ there are 2-submanifolds of relative equilibria, as α_e is varied, as expected for an $SO(2)$ symmetric Hamiltonian system. Along those submanifolds there is the *momentum-generator* relation

$$\hat{\xi}_e^{\alpha_e} = \frac{1}{I_3} \hat{\mu}_e^{\alpha_e}, \quad (7)$$

which will give a crucial component in the KAM twist condition to follow.

§2 NORMAL FORMS IN THE BLOWN-UP SYSTEM

Proposition 1 below relates the stability of the relative equilibrium $p_e^{\alpha_e}$ to the stability of its blow-up $\hat{p}_e^{\alpha_e}$.

Proposition 1 *Suppose that, for some fixed α_e , $\hat{p}_e^{\alpha_e}$ is stable for the flow $\hat{F}_t$ on $\hat{P}$, in the sense that for all neighborhoods $\hat{U}$ of $\hat{p}_e^{\alpha_e}$ there is a neighborhood $\hat{V}$ of $\hat{p}_e^{\alpha_e}$ such that $\hat{F}_t(\hat{p}) \in \hat{U}$ for all $\hat{p} \in \hat{V}$. Then $p_e^{\alpha_e}$ is a stable relative equilibrium.*

Proof. Suppose U is a neighborhood of $p_e^{\alpha_e}$. U pulls back by the blow-down map to an open neighborhood $\hat{U}$ of $\hat{p}_e^{\alpha_e}$. Let $\hat{V}$ be a neighborhood as in the statement of the proposition. Then it suffices to show that $\hat{V}$ pushes forward by the blow-down map to a neighborhood of $p_e^{\alpha_e}$. But this follows since the blow-down map is proper. ■

In particular, if all of the relative equilibria and both equilibria in $\hat{p}_e^{\alpha_e}$ are stable under perturbation both within the phase space TS^2 and in the parameters a and γ , then the original relative equilibrium $p_e^{\alpha_e}$ is stable. When a is perturbed away from 0 this is a $SO(2)$ symmetry breaking perturbation. As $\dim TS^2 = 4$, the blown-up system is integrable when $a = 0$ and hence the stability issue is one of the stability of periodic orbits of a near integrable Hamiltonian system.

Assume, without loss of generality, that $\alpha_e > 0$. Since the $SO(2)$ symmetry on $\hat{p}_e^{\alpha_e}$ is by rotation about $\mathbf{k}$, it suffices to consider the stability of orbits in $\hat{p}_e^{\alpha_e}$ emanating from points $\hat{p}_e^{\alpha_e, \theta}$ obtained by substituting $w = \sin\theta\mathbf{i} + \cos\theta\mathbf{k}$ into (5), for $\theta \in [0, \pi]$.

§2.1 NORMAL FORM FOR THE RELATIVE EQUILIBRIA

Consider first the *relative equilibria* in $\hat{p}_e^{\alpha_e, \theta}$; i.e., exclude the *equilibria* corresponding to $\theta = 0$ and $\theta = \pi$. The argument proceeds by adapting and incrementally refining, to the order required for the stability analysis, the normal form near relative equilibria developed in Patrick [1995].

Below $O(x; y)^k$, $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$ will denote the set of smooth y dependent functions such that $O(x; y)/|x|^k$ is bounded near 0. The product $O(x; y)^k O(x'; y')^{k'}$ denotes the set of finite sums of products of elements of $O(x; y)^k$ and $O(x'; y')^{k'}$.

§2.1.1 *Initial normal form.* This is constructed from the linearization of the relative equilibrium, which means the linearization at $\hat{p}_e^{\alpha_e, \theta}$ of Hamiltonian vector field $X_{\hat{H}_{\xi_e^{\alpha_e}}}$ where

$$\hat{H}_{\xi_e^{\alpha_e}}^0 \equiv \hat{H}^0 - \hat{\xi}_e \hat{J}.$$

The characteristic polynomial of the linearization is $x \mapsto x^2(x^2 + \omega_e^2)$, where

$$\omega_e \equiv \pm \alpha_e \sqrt{\left(\frac{1}{I_3} - \frac{1}{I_1}\right) \left(\frac{1}{I_3} - \frac{1}{I_2}\right)}, \quad (8)$$

For later convenience define ω_e positive if $I_3 > I_1$ and $I_3 > I_2$ and negative if $I_3 < I_1$ and $I_3 < I_2$. The linearization has a 0 and $\pm i\omega_e$ generalized eigenspaces, both of dimension 2. Introducing the parameter

$$D \equiv \frac{I_2(I_3 - I_1)}{I_1(I_3 - I_2)},$$

the vectors

$$\begin{aligned} v_1 &\equiv \frac{D^{\frac{1}{4}}}{\sqrt{\alpha_e}} \begin{bmatrix} 0 & \cos \theta & 0 & 0 & \alpha_e \sin^2 \theta & 0 \end{bmatrix} \\ v_2 &\equiv \frac{D^{-\frac{1}{4}}}{\sqrt{\alpha_e}} \begin{bmatrix} \cos \theta & 0 & -\sin \theta & \alpha_e \sin^2 \theta & 0 & \alpha_e \sin \theta \cos \theta \end{bmatrix} \\ v_3 &\equiv \sin \theta \begin{bmatrix} 0 & 1 & 0 & 0 & -\alpha_e \cos \theta & 0 \end{bmatrix} \\ v_4 &\equiv \frac{1}{\alpha_e \sin \theta} \begin{bmatrix} -\cos^2 \theta & 0 & \cos \theta \sin \theta & \alpha_e \cos^3 \theta & 0 & -\alpha_e \sin \theta (1 + \cos^2 \theta) \end{bmatrix} \end{aligned}$$

form a basis of $T_{\hat{p}_e^{\alpha_e, \theta}} S^2$ which satisfies the following:

1. the basis is symplectically canonical, so that the symplectic form with respect to it is

$$\omega(\hat{p}_e^{\alpha_e, \theta}) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix};$$

2. with respect to the basis the derivative of the momentum is

$$d\hat{J}(\hat{p}_e^{\alpha_e, \theta}) = \begin{bmatrix} 0 & 0 & 0 & 1 \end{bmatrix};$$

3. the third basis vector v_3 is the infinitesimal generator action corresponding to $1 \in so(2)$;

4. the first two vectors v_1, v_2 span the ω_e generalized eigenspace and the last two v_3, v_4 span the 0 generalized eigenspace of the linearization;
5. the linearization of the relative equilibrium is

$$dX_{\hat{H}_{\xi_e^{\alpha_e}}^0}(\hat{p}_e^{\alpha_e, \theta}) = \begin{bmatrix} 0 & \omega_e & 0 & 0 \\ -\omega_e & 0 & 0 & 0 \\ 0 & 0 & 0 & \kappa_e \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (9)$$

where

$$\kappa_e \equiv 1/I_3. \quad (10)$$

Consequently, the basis effects a Witt-Moncrief decomposition

$$T_{\hat{p}_e^{\alpha_e, \theta}} \hat{P} = N_1 \oplus so(2) \oplus so(2)^* \equiv \text{span}(v_1, v_2) \oplus \mathbb{R}v_3 \oplus \mathbb{R}v_4.$$

Here N_1 , the *symplectic normal*, may be identified with the tangent space to the symplectic reduced space at $\hat{p}_e^{\alpha_e, \theta}$ for the $SO(2)$ action. The appearance of the nilpotent part of the linearization is the foundational element of Patrick [1995]. The value of κ_e coincides with the derivative $d\hat{\xi}_e^{\alpha_e}/d\hat{\mu}_e^{\alpha_e}$ from (7), as predicted by the general theory.

The initial normal form can now transcribed from the data above, and is

$$\begin{aligned} \hat{H} &= \frac{\omega_e}{2}(q^2 + p^2) + \hat{\xi}_e^{\alpha_e} \nu + \frac{1}{2} \kappa_e \nu^2 + R(q, p, \nu) + \frac{a^2}{2} \hat{H}^1(q, p, \varphi, \nu), \\ R &= O(q, p, \nu)^3 \end{aligned} \quad (11)$$

on the product of $\mathbb{R}^2 \times T^*SO(2) = \{(q, p), (\varphi, \nu)\}$ with the product symplectic form $dq \wedge dp + d\varphi \wedge d\nu$, with $SO(2)$ acting by lifts of its left action on itself, and with the momentum map $\nu - \alpha_e$. The transcription is that there is an $SO(2)$ intertwining symplectic diffeomorphism from a neighborhood of the $SO(2)$ orbit of $\hat{p}_e^{\alpha_e}$ to a neighborhood of 0 times the zero section of $T^*SO(2)$ which

1. sends the relative equilibrium $\hat{p}_e^{\alpha_e, \theta}$ to $p = q = \nu = \varphi = 0$.
2. has derivative at $\hat{p}_e^{\alpha_e}$ the identity map with respect to the basis v_i and the standard basis of $\mathbb{R}^2 \times T^*SO(2)$;
3. intertwines the momentum maps $\hat{J}$ and $\nu - \alpha_e$;

Thus the transription is *structure preserving* in that it is symplectic and it preserves the $SO(2)$ symmetry and momentum, so the blown-up system near the group orbit of the relative equilibrium $\hat{p}_e^{\alpha_e, \theta}$ can be replaced by the entirely equivalent system (11) near $q = p = \nu = 0$.

§2.1.2 *Elimination of $qO(\nu)^2$, $pO(\nu)^2$, and $(q^2 - p^2)\nu$.* The remainder term of (11) can be expanded as

$$R = c_1(\nu)q + c_2(\nu)p + c_3\nu(q^2 - p^2) + c_4\nu(q^2 + p^2) + O(q, p)^3 + O(\nu)^3 + O(q, p)^2 O(q, p, \nu)^2,$$

where $c_1(\nu) = O(\nu)^2$, $c_2(\nu) = O(\nu)^2$, and c_3, c_4 are constants. The transformation

$$\tilde{q} = q + \frac{c_1}{\omega_e}, \quad \tilde{\varphi} = \varphi + \frac{p\nu}{\omega_e} \frac{dc_1}{d\nu},$$

suggested by completing the square in $\frac{1}{2}\omega_e q^2 + c_1 q \nu^2$, is structure preserving and changes the Hamiltonian to the same form but without terms of the form $qO(\nu)^2$. Similarly one eliminates $pO(\nu)^2$. The transformation

$$\tilde{q} = \frac{q}{f(\nu)}, \quad \tilde{p} = f(\nu)p, \quad f(\nu) = \left(\frac{1 - \frac{2c_3}{\omega_e}\nu}{1 + \frac{2c_3}{\omega_e}\nu} \right)^{\frac{1}{4}}, \quad (12)$$

takes the fragment $\frac{\omega_e}{2}(q^2 + p^2) + c_3(q^2 - p^2)\nu$ to

$$\left(\frac{\omega_e}{2} + c_3 \right) q^2 + \left(\frac{\omega_e}{2} - c_3 \right) p^2 = \frac{\omega_e}{2}(\tilde{q}^2 + \tilde{p}^2) + \tilde{q}^2 O(\nu)^2 + \tilde{p}^2 O(\nu)^2,$$

while the symplectic form becomes

$$\begin{aligned} dq \wedge dp + d\varphi \wedge d\nu &= d\tilde{q} \wedge dp + d\varphi \wedge d\nu + \frac{f'}{f}(qdp + pdq) \wedge d\nu \\ &= d\tilde{q} \wedge dp + d \left(\varphi + \frac{f'}{f}qp \right) \wedge d\nu. \end{aligned}$$

Adjoining $\tilde{\varphi} = \varphi + (f'/f)qp$ to (12) gives a structure preserving symplectic transformation that eliminates the term $c_3(q^2 - p^2)$. Thus, without loss of generality,

$$R = c_4(q^2 + p^2) + O(q, p)^3 + O(\nu)^3 + O(q, p)^2 O(q, p, \nu)^2. \quad (13)$$

§2.1.3 *Normal form for the rigid body.* We will require the first two terms of the normal form corresponding to the equilibrium $\pi = \alpha_e \mathbf{k}$ of the blown-up system reduced by its $SO(2)$ symmetry, i.e. the symplectic reduced spaces of the rigid body $\frac{1}{2}\pi \mathbf{I}^{-1} \pi$. The map

$$\pi = \left(\left(\alpha_e - \frac{1}{4}(Q^2 + P^2) \right)^{\frac{1}{2}} P, \left(\alpha_e - \frac{1}{4}(Q^2 + P^2) \right)^{\frac{1}{2}} Q, \alpha_e - \frac{1}{2}(Q^2 + P^2) \right)$$

is a symplectic chart on the reduced space $|\pi| = \alpha_e$ and in these coordinates, the Hamiltonian becomes, up to a constant,

$$\frac{1}{2}\pi^t \mathbf{I}^{-1} \pi = \frac{1}{2} \left(\alpha_e - \frac{1}{4}(Q^2 + P^2) \right) \left(\left(\frac{1}{I_1} - \frac{1}{I_3} \right) P^2 + \left(\frac{1}{I_2} - \frac{1}{I_3} \right) Q^2 \right).$$

Action-angle variables for the linearized flow are

$$Q = \sqrt{2I}D^{\frac{1}{4}} \sin \psi, \quad P = \sqrt{2I}D^{-\frac{1}{4}} \cos \psi,$$

and the Hamiltonian is then

$$\frac{1}{2}\pi^t \mathbf{I}^{-1} \pi = \omega_e I - \frac{\omega_e}{2\alpha_e} (D^{\frac{1}{2}} \sin^2 \psi + D^{-\frac{1}{2}} \cos^2 \psi) I^2.$$

By averaging over ψ ,

$$\frac{1}{2}\pi^t \mathbf{I} \pi = \omega_e I + \frac{1}{2}v_e I^2 + O(Q, P)^5 \quad (14)$$

where

$$v_e = -\frac{\omega_e}{2\alpha_e} (D^{\frac{1}{2}} + D^{-\frac{1}{2}}) = \frac{1}{2} \left(\frac{2}{I_3} - \frac{1}{I_1} - \frac{1}{I_2} \right). \quad (15)$$

§2.1.4 Matching and normalizing the reduced spaces at $\hat{p}_e^{\alpha_e, \theta}$. For $a = 0$ the symplectic reduced space through $\hat{p}_e^{\alpha_e, \theta}$ of the blown-up system is the $|\pi| = \alpha_e$ symplectic reduced space of the rigid body $\frac{1}{2}\pi^t \mathbf{I} \pi$. For $a = 0$ the symplectic reduced space of the normal form (11) through $q = p = \varphi = \nu = 0$ is $\mathbb{R}^2$ with symplectic form $dq \wedge dp$ and Hamiltonian $\hat{H}|_{\nu=0}$. Since the intertwining map between the blown-up system and the normal form is structure preserving, it descends to symplectomorphisms of reductions of these two systems. Consequently, by symplectomorphism on (q, p) only, the normal form Hamiltonian (11) at $q = p = \nu = 0$ can be equated to the rigid body normal form (14), after which the normal is correct to fourth order in pure q and p and

$$R = O(q, p)^5 + \nu O(q, p)^2 O(q, p, \nu).$$

§2.1.5 Refinement by matching the normal forms and generators along the relative equilibria near $\hat{p}_e^{\alpha_e, \theta}$. Advantage may be obtained by comparing reduced normal along the relative equilibria $\hat{p}_e^{\alpha_e + z, \theta}$ as z varies. These relative equilibria occur (for both systems) at momentum $-(\alpha_e + z)$. For the rigid body the only z dependent adjustment is in the α_e dependence of the linearized frequency, which becomes $\omega_e(\alpha_e + z)/\alpha_e$, so the normal form is

$$\left(\omega_e + \frac{\omega_e}{\alpha_e} z \right) I + \frac{1}{2}v_e I^2 + O(q, p; z)^5. \quad (16)$$

For (11) it is the normal form of the reduction at $\nu = -z$, so it is the normal form of the Hamiltonian

$$\omega_e I + \frac{1}{2}v_e I^2 - 2c_4 I z + O(q, p)^2 O(q, p, z)^2$$

which is

$$(\omega_e - 2c_4 z + O(z)^2) I + O(q, p; z)^3. \quad (17)$$

Comparison of (16) and (17) at first order in I gives a crucial fact:

$$c_4 = -\frac{\omega_e}{2\alpha_e}.$$

Also, the $SO(2)$ generator of the blown-up system at $\hat{p}_e^{\alpha_e+z, \theta}$, which is $-(\alpha_e + z)/I_3$, and the $SO(2)$ generator of system (11) at the relative equilibrium $q = p = 0$ are the same. Equating these gives

$$\hat{\xi}_e^{\alpha_e} + \kappa_e \nu + \left. \frac{\partial R}{\partial \nu} \right|_{\substack{q=p=0 \\ \nu=-z}} = -\frac{1}{I_3}(\alpha_e + z) = \hat{\xi}_e^{\alpha_e} - \kappa_e z,$$

which means that R has no pure ν terms. Particularly, the $O(\nu)^3$ term in (13) is zero.

§2.1.6 Symmetry breaking term. The transcription to the initial normal form is known to first order since it has derivative the identity map along the $SO(2)$ orbit of $\hat{p}_e^{\alpha_e, \theta}$. Consequently, $\hat{H}^1$ can be calculated to first order by substitution of

$$w = \exp(\varphi \mathbf{k}) P_w(\hat{p}_e^{\alpha_e, \theta} + (qv_1 + pv_2 + \nu v_4))$$

into $w^t \mathbf{M}^{-1} w$, where $P_w(w, \dot{w}) = w$.

§2.1.7 Altogether. Putting all the foregoing together, the normal form is

$$\begin{aligned} \hat{H} = & \omega_e I + \frac{1}{2} v_e I^2 + \hat{\xi}_e^{\alpha_e} \nu + \frac{1}{2} \kappa_e \nu^2 - \frac{\omega_e}{\alpha_e} I \nu \\ & + O(q, p)^5 + \nu O(q, p)^2 O(q, p, \nu) \\ & + a^2 \hat{H}^{1,0}(q, p, \varphi, \nu) + a^2 \hat{H}^{1,1}(q, p, \varphi, \nu) + a^2 O(q, p, \nu; \varphi)^2, \end{aligned} \quad (18)$$

where

$$\hat{H}^{1,0} \equiv \frac{M_2 - M_1}{2M_1 M_2} \sin^2 \theta \cos^2 \varphi \quad (19)$$

and

$$\begin{aligned} \hat{H}^{1,1} \equiv & \frac{(M_2 - M_1) \sin 2\theta}{4M_1 M_2 \sqrt{\alpha_e}} \left(-D^{\frac{1}{4}} q \sin 2\varphi + D^{-\frac{1}{4}} p \cos 2\varphi \right) \\ & - \frac{\cos^2 \theta}{\alpha_e} \left(\frac{1}{M_1} \cos^2 \varphi + \frac{1}{M_2} \sin^2 \varphi - \frac{1}{M_3} \right) \nu \\ & - \frac{\sin 2\theta}{4D^{\frac{1}{4}} \sqrt{\alpha_e}} \left(\frac{2}{M_3} - \frac{1}{M_1} - \frac{1}{M_2} \right) p. \end{aligned}$$

The details of the symmetry breaking term $\hat{H}^1$ are not required for the stability analysis and are displayed here for the sake of completeness. The functional form of $\hat{H}^{1,1}$ depends on the choice of the basis v_i and further normalization or analysis would be required to extract information from it.

§2.2 NORMAL FORM FOR THE EQUILIBRIA

There remains to consider the two equilibria $\hat{p}_e^{\alpha_e,0}$ and $\hat{p}_e^{\alpha_e,\pi}$ corresponding to $w = \mathbf{k}$ and $w = -\mathbf{k}$, respectively. These equilibria are fixed points of the action of $SO(2)$ and the analysis requires a transparent extension of the normal form in Patrick [1995] to *equilibria* which have $SO(2)$ isotropy.

It suffices to consider $\hat{p}_e^{\alpha_e,0}$; the case of $\hat{p}_e^{\alpha_e,\pi}$ is similar. There is a one parameter family of possible linearizations of the equilibrium, namely the linearizations at $\hat{p}_e^{\alpha_e,0}$ of the Hamiltonian vector fields $X_{\hat{H}_\lambda^0}$ where $\hat{H}_\lambda^0 = \lambda \hat{J}$. These linearizations have characteristic polynomials

$$x \mapsto (x^2 + (\alpha_e + \lambda I_3)^2)(x^2 + \omega_e^2).$$

Choosing $\lambda = -\alpha_e/I_3$ gives the largest possible null space and therefore the largest number of intrinsically defined higher order terms. The vectors

$$\begin{aligned} v_{1,0} &\equiv \frac{D^{\frac{1}{4}}}{\sqrt{\alpha_e}} \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \end{bmatrix}, & v_{2,0} &\equiv \frac{D^{-\frac{1}{4}}}{\sqrt{\alpha_e}} \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \\ v_{3,0} &\equiv \frac{1}{\sqrt{\alpha_e}} \begin{bmatrix} 0 & 1 & 0 & 0 & -\alpha_e & 0 \end{bmatrix} & v_{4,0} &\equiv \frac{1}{\sqrt{\alpha_e}} \begin{bmatrix} -1 & 0 & 0 & \alpha_e & 0 & 0 \end{bmatrix}, \end{aligned}$$

form a basis of $T_{\hat{p}_e^{\alpha_e,0}} S^2$ which is symplectically canonical and such that the linearization $dX_{H_\lambda^0}(\hat{p}_e^{\alpha_e,0})$ is the linearization (9) with κ_e replaced by zero. The first two vectors span the tangent space to the (singular) reduced space through $p_e^{\alpha_e,0}$, which is the $|\pi| = \alpha_e$ symplectic reduced space of the rigid body $\frac{1}{2}\pi^t \mathbf{I}^{-1} \pi$.

Higher order terms of the Taylor expansion of $X_{H_\lambda^0}$ are intrinsically polynomials on $\ker dX_{H_\lambda^0}(\hat{p}_e^{\alpha_e,0})$. Alternately one can compute the higher order terms of the Taylor expansion of the Hamiltonian on the null space. Letting (x, y) be the coordinates on $\ker dX_{H_\lambda^0}(\hat{p}_e^{\alpha_e,0})$ indicated by the last of the two basis vectors above, the Hamiltonian on the null space is easily computed to be

$$\frac{1}{8I_3}(x^2 + y^2)^2 + O(x, y)^5 = \frac{\kappa_e}{8}(x^2 + y^2)^2 + O(x, y)^5.$$

The initial normal form, obtained by the Equivariant Darboux Theorem, is the Hamiltonian

$$\begin{aligned} \hat{H} &= \frac{\omega_e}{2}(q^2 + p^2) - \frac{\alpha_e}{2I_3}(x^2 + y^2) + \frac{\kappa_e}{8}(x^2 + y^2)^2 \\ &\quad + O(q, p)O(q, p, x, y)^2 + O(x, y)^5 + a^2 \hat{H}^1(q, p, x, y) \end{aligned}$$

on the phase space $\mathbb{R}^2 \times \mathbb{R}^2 = \{(q, p), (x, y)\}$, with symplectic form $dq \wedge dp + dx \wedge dy$, with $SO(2)$ acting by counterclockwise rotation on (x, y) , and with the momentum mapping $-\frac{1}{2}(x^2 + y^2)$. The transcription is by local symplectic diffeomorphism with analogous properties to those stated in Section 2.1.1.

Manipulations similar to those in Section 2.1 are required, as follows:

1. Linear terms in q of the form $qO(x, y)^2$ can be removed as in Section 2.1.2, and similarly linear terms in p , x , and y . Quartic terms of the form $O(q, p)^2O(x, y)^2$ must by $SO(2)$ invariance be in $(x^2 + y^2)O(q, p)$, and so can be written as sums of $(q^2 + p^2)(x^2 + y^2)$ and $(q^2 - p^2)(x^2 + y^2)$, and the latter kind removed, as in Section 2.1.2.
2. Pure q and p terms up to order 4 can be found by matching the reduced system of the initial normal form to rigid body reduced spaces.
3. By Items 1 and 2 all terms up to and including order 4 are removed or calculated, except for the coefficient of the term $(q^2 + p^2)(x^2 + y^2)$. This can be found by matching normal forms along the equilibria $q = p = 0$ (which are fixed points of the action of $SO(2)$), and the resulting term is $-\frac{\omega_e}{4\alpha_e}(q^2 + p^2)(x^2 + y^2)$.
4. The remainder after all of that, having no terms linear in any variable, and being at least degree 5, is of the form $O(q, p, x, y)^2O(q, p, x, y)^3$, and is $SO(2)$ invariant.
5. The symmetry breaking term $\hat{H}^1$ can be calculated as in Section 2.1.6 by substituting

$$w = P_w(\hat{p}_e^{\alpha_e, 0} + qv_{1,0} + pv_{2,0} + xv_{3,0} + yv_{4,0})$$

into $w^t \mathbf{M}^{-1}w$, and keeping the leading terms, which are order 2.

Altogether, the normal form is

$$\begin{aligned} \hat{H} = & \omega_e I + \frac{1}{2}v_e I^2 + \xi_e^{\alpha_e} \nu + \frac{1}{2}\kappa_e \nu^2 - \frac{\omega_e}{\alpha_e} I \nu + O(q, p, x, y)^2 O(q, p, x, y)^3 \\ & + \frac{a^2}{2} \hat{H}^{1,1}(q, p, x, y) + a^2 O(q, p, x, y)^3 \end{aligned} \quad (20)$$

where $I = \frac{1}{2}(q^2 + p^2)$ and $\nu = \frac{1}{2}(x^2 + y^2)$, and

$$\hat{H}^{1,1} \equiv \frac{(M_3 - M_1)(D^{-\frac{1}{4}}p - y)^2}{\alpha_e M_1 M_3} + \frac{(M_3 - M_2)(D^{\frac{1}{4}}q + x)^2}{\alpha_e M_2 M_3}.$$

§3 STABILITY

The rescaling

$$I = a^{2c} \tilde{I}, \quad \nu = a^{2c} \tilde{\nu}$$

is symplectic with multiplier a^{2c} . Substituting into (18) and dropping the tildes gives

$$\begin{aligned} \frac{1}{a^{2c}} \hat{H} = & \omega_e I + \frac{1}{2}v_e I^2 a^{2c} + \xi_e^{\alpha_e} \nu + \frac{1}{2}\kappa_e \nu^2 a^{2c} - \omega_e I \nu a^{2c} + O(a; q, p, \varphi, \nu)^{3c} \\ & + a^{2-2c} \hat{H}^{1,0} + O(a; q, p, \varphi, \nu)^{2-c} \end{aligned}$$

Matching the exponents of a in first nontrivial terms of the integrable part, i.e. $\frac{1}{2}\kappa_e\nu^2a^{2c}$ and $\frac{1}{2}v_eI^2a^{2c}$, with the first term of the nonintegrable part, gives $2c = 2 - 2c$, or $c = \frac{1}{2}$. After putting $\epsilon = \sqrt{a}$, and disposing the factor $1/a^{2c}$ of $\hat{H}$, which merely reparameterizes time, one has

$$\hat{H} = \omega_e I + \hat{\xi}_e^{\alpha_e} \nu + \left(\frac{1}{2} \kappa_e \nu^2 - \frac{\omega_e}{\alpha_e} I \nu + \frac{1}{2} v_e I^2 + \hat{H}^{1,0} \right) \epsilon^2 + O(\epsilon)^3, \quad (21)$$

where the dependence of $O(\epsilon)^3$ on all of q , p , φ , and ν has been notationally suppressed. For $\epsilon = 0$ the Hamiltonian (21) has a periodic orbit cylinder by varying φ and ν with $I = 0$. The orbit $I = \nu = 0$ corresponds to the relative equilibrium $\hat{p}_e^{\alpha_e, \theta}$.

For determining stability it suffices to approximate the Poincaré map for the orbit corresponding to $\nu = 0$ in the zero energy level. Solving (21) for ν when $H = 0$ gives

$$\nu = -\frac{\omega_e I}{\hat{\xi}_e^{\alpha_e}} + O(\epsilon)^2$$

and the equations of motion for (21) are

$$\begin{aligned} \frac{d\psi}{dt} &= \omega_e + \left(v_e I - \frac{\omega_e}{\alpha_e} \nu \right) \epsilon^2 + O(\epsilon)^3, & \frac{dI}{dt} &= O(\epsilon)^3, \\ \frac{d\varphi}{dt} &= \hat{\xi}_e^{\alpha_e} + \left(\kappa_e \nu - \frac{\omega_e}{\alpha_e} I \right) \epsilon^2 + O(\epsilon)^3, & \frac{d\nu}{dt} &= -\frac{\partial \hat{H}^{1,0}}{\partial \varphi} \epsilon^2 + O(\epsilon)^3. \end{aligned} \quad (22)$$

On the zero energy level the equation for the evolution of φ is

$$\frac{d\varphi}{dt} = \hat{\xi}_e^{\alpha_e} - \frac{\omega_e(\kappa_e \alpha_e + \hat{\xi}_e^{\alpha_e})}{\alpha_e \hat{\xi}_e^{\alpha_e}} I \epsilon^2 + O(\epsilon)^3. \quad (23)$$

With an initial condition $I = I_0$, the second equation of (22) gives $I = I_0 + O(\epsilon)^3$, so the return time of the Poincaré map is, from (23),

$$T \equiv -\frac{2\pi}{\hat{\xi}_e^{\alpha_e}} \left(1 + \frac{(\alpha_e \kappa_e + \hat{\xi}_e^{\alpha_e}) \omega_e I_0}{\alpha_e (\hat{\xi}_e^{\alpha_e})^2} \epsilon^2 \right) + O(\epsilon)^3$$

Solving the first two equations of (22) over this period, dropping the subscript 0 for the initial conditions, and using the relation $\hat{\xi}_e^{\alpha_e} = -\kappa_e \alpha_e$ to eliminate α_e gives

$$\begin{aligned} I' &= I + O(\epsilon)^3 \\ \psi' &= \psi - \frac{2\pi \omega_e}{\hat{\xi}_e^{\alpha_e}} - \epsilon^2 \frac{2\pi}{(\hat{\xi}_e^{\alpha_e})^3} (v_e (\hat{\xi}_e^{\alpha_e})^2 - \kappa_e \omega_e^2) I + O(\epsilon)^3 \end{aligned} \quad (24)$$

This is of the form (Meyer and Hall [1991], Theorem 2, page 231), namely $(I, \psi) \mapsto (I', \psi')$ by

$$\begin{aligned} I' &= I + \epsilon^{r+s} c(I, \psi, \epsilon), \\ \psi' &= \psi + \omega + \epsilon^s h(I) + \epsilon^{s+r} d(I, \psi, \epsilon) \end{aligned}$$




Figure 1: Numerical verification of the twist predicted by (24). Leftmost: the curved leading edge of the numerically computed Poincaré map indicates a twist map by visibly showing faster rotation as I increases. The twist overlays a constant rotation (in I) which is caused by high order terms in a and decreases as a decreases. Center: the rotation angle per iteration of the Poincaré map on the right as a function of I . The slope corresponds to the twist predicted by (24). Right: reciprocal of the twist for $I_3 = 1$, $I_2 = I_3 - I_1$ as I_1 ranges from 0 to 1 compared to the parabola predicted by (24).

with $r = 1$, $s = 2$, and

$$h(I) \equiv -\frac{2\pi}{(\hat{\xi}_e^{\alpha_e})^3} (v_e(\hat{\xi}_e^{\alpha_e})^2 - \kappa_e \omega_e^2) I.$$

The twist condition $dh/dI \neq 0$ is

$$v_e(\hat{\xi}_e^{\alpha_e})^2 - \kappa_e \omega_e^2 \neq 0, \quad (25)$$

which, after substituting (6), (8), (10), and (15), is

$$\begin{aligned} & -\frac{1}{2} \left(\frac{2}{I_3} - \frac{1}{I_1} - \frac{1}{I_2} \right) + I_3 \left(\frac{1}{I_3} - \frac{1}{I_1} \right) \left(\frac{1}{I_3} - \frac{1}{I_2} \right) \\ & = \frac{1}{I_1 I_2} \left(I_3 - \frac{1}{2}(I_1 + I_2) \right) \neq 0. \end{aligned}$$

This is certainly true if I_3 is not between I_1 and I_2 .

As for the equilibria $\hat{p}_e^{\alpha_e, 0}$ of Section 2.2, Arnold's Stability Theorem (Meyer and Hall [1991], Theorem 1, page 235) together with the normal form (20) imply that the equilibrium is stable when $a = 0$ if

$$\omega_e I + \frac{1}{2} v_e I^2 + \hat{\xi}_e^{\alpha_e} \nu + \frac{1}{2} \kappa_e \nu^2 - \frac{\omega_e}{\alpha_e} I \nu \Big|_{\substack{I = \hat{\xi}_e^{\alpha_e} \\ \nu = -\omega_e}} = \frac{1}{2} (v_e(\hat{\xi}_e^{\alpha_e})^2 - \kappa_e \omega_e^2) \neq 0,$$

which is the same as the twist condition (25). Stability follows for sufficiently small nonzero a since a contributes continuously.

Thus $p_e^{\alpha_e}$ are stable as equilibria on the Poisson reduced space $\{(\pi, p)\}$. From Patrick, Roberts, and Wulff [2002], and since the momentum at $p_e^{\alpha_e}$ has zero translational part and rotational part parallel to $\mathbf{k}$, this implies $SO(2) \times C$ stability of the original equilibrium, where C is any cone about $\mathbf{k}$.

The foregoing sort of analysis will always lead to some stability condition, irrespective of possible errors in the derivation, so it necessary to check (24) by

comparing it with numerically generated Poincaré maps. Substitution of $\epsilon = 1$ after truncation of $O(\epsilon)^3$ into (24) gives the leading behavior of the Poincaré map when a , ν , and I , are of comparable order (they are all order ϵ^2). The Poincaré map is determined to leading order in I by the first order twist term $h(I)$ in when the zero order term $2\pi\omega_e/\hat{\xi}_e^{\alpha_e}$ has a vanishing effect (i.e. is a multiple of 2π). This happens to occur when $I_1 + I_2 = I_3$, as is easily verified. After substitution of $I_2 = I_3 - I_1$, the first order twist is

$$h(I) = -\frac{\pi I_3^2}{I_1(I_3 - I_1)}I,$$

whereupon

$$\frac{I}{h(I)} = -\frac{1}{\pi I_3^2}I_1(I_3 - I_1),$$

which is a parabola in I_1/I_3 . As can be seen in Figure 1, this compares well with numerical integrations of the original (as opposed to the blown-up) system.

SUMMARY

The following theorem has been proved.

Theorem 1 *Within the context of the Lagrangian system (1), the motion of an underwater ellipsoid rotating about a long or short principle axis of inertia is stable modulo $SO(2) \times C$ where $SO(2)$ acts around the rotation axis and C is any cone containing that axis.*

This theorem follows from KAM confinement after a blow-up construction and normal form analysis rather than confinement by Lyapunov functions derived from energy and momentum.

ACKNOWLEDGMENTS

This work was supported by an NSERC individual research grant and an EPSRC visiting fellowship. I thank the University of Warwick Mathematics Institute for its hospitality during a sabbatical visit while this paper was written.

REFERENCES

- Leonard, N. E. [1997]. Stability of a bottom-heavy underwater vehicle. *Automatica—J. IFAC* 33, 331–346.
- Leonard, N. E. and J. E. Marsden [1997]. Stability and drift of underwater vehicle dynamics: mechanical systems with rigid motion symmetry. *Physica D* 105, 130–162.
- Marsden, J. E. and T. S. Ratiu [1994]. *Introduction to Mechanics and Symmetry*, Volume 17 of *Texts in Applied Mathematics*. Springer-Verlag.
- Meyer, K. R. and G. R. Hall [1991]. *Introduction to Hamiltonian dynamical systems and the N-body problem*. Springer-Verlag.

- Patrick, G., R. M. Roberts, and C. Wulff [2002]. Stability of Poisson equilibria and Hamiltonian relative equilibria by energy methods. Preprint.
- Patrick, G. W. [1995]. Relative equilibria of Hamiltonian systems with symmetry: linearization, smoothness, and drift. *J. Nonlinear Sci.* 5, 373–418.